

EVOLUTION OF THE IR ENERGY DENSITY AND SFH UP TO $Z \sim 1$: FIRST RESULTS FROM MIPS

Emeric Le Floch¹, C. Papovich¹, H. Dole², E. Egami¹, P. Pérez-González¹, G. Rieke¹, M. Rieke¹, E. Bell³ & the *Spitzer*/MIPS GTO team¹

¹ *Steward Observatory, University of Arizona, 933, N. Cherry Avenue, Tucson 85721, AZ, USA*

² *Institut d'Astrophysique Spatiale, Université Paris Sud, F-91405 Orsay Cedex, France*

³ *Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

(elefloch@as.arizona.edu)

Abstract Using deep observations of the Chandra Deep Field South obtained with MIPS at $24 \mu\text{m}$, we present our preliminary estimates on the evolution of the infrared (IR) luminosity density of the Universe from $z=0$ to $z \sim 1$. We find that a pure density evolution of the IR luminosity function is clearly excluded by the data. The characteristic luminosity L_{IR}^* evolves at least by $(1+z)^{3.5}$ with lookback time, but our monochromatic approach does not allow us to break the degeneracy between a pure evolution in luminosity or an evolution in both density and luminosity. Our results imply that IR luminous systems ($L_{\text{IR}} \geq 10^{11} L_{\odot}$) become the dominant population contributing to the comoving IR energy density beyond $z \sim 0.5-0.6$. The uncertainties affecting our measurements are largely dominated by the poor constraints on the spectral energy distributions that are used to translate the observed $24 \mu\text{m}$ flux into luminosities.

Keywords: Galaxy evolution, observational cosmology, infrared luminous galaxies

Introduction

Deep infrared and submillimeter observations performed with *ISO* and SCUBA in the late 90s revealed a very strong evolution of IR luminous systems with lookback time (e.g., Smail et al. 1997, Elbaz et al. 1999). Since these surveys were only sensitive to the brightest IR sources, the quantification of this evolution has yet remained strongly debated so far. In particular, the relative importance of such IR-bright objects compared to less luminous starbursts, and their respective contributions to the total comoving energy density at high redshift is still unclear. *Spitzer*, the new infrared facility of NASA, is now providing a unique opportunity to address this issue in more detail.

1. Infrared luminosities of MIPS sources up to $z \sim 1$

We performed deep observations of the Chandra Deep Field South with MIPS at $24 \mu\text{m}$ down to a sensitivity limit of $80 \mu\text{Jy}$ (5σ detection). Cross-correlating our data with source catalogs from various optical surveys (i.e., VIMOS VLT Deep Survey: Le Fèvre et al. 2004; GOODS: Vanzella et al. 2004; COMBO-17: Wolf et al. 2004; Chandra source follow-up: Szokoly et al. 2004), we derived the redshift of 2635 objects detected at $24 \mu\text{m}$ and mostly located at $z \leq 1.2$. We believe that this redshift identification is nearly complete up to $z \sim 1$.

Using various libraries of IR luminosity-dependent spectral energy distributions (SEDs, e.g., Dale et al. 2001, Chary & Elbaz 2001), we derived the total IR luminosities of the MIPS sources from their observed $24 \mu\text{m}$ flux. Given the poor constraint on the true SEDs characterizing these sources at high redshift, the typical uncertainty affecting these estimates could reach a factor of 2 to 3. More importantly, influence of dust temperature could also add a small systematic bias (Chapman et al. 2003) *that we have not quantified so far*. We will explore this effect more thoroughly in a forthcoming work.

Below $z \sim 0.5$, we find that the MIPS sources are rather modest emitters at infrared wavelengths (median $L_{\text{IR}} = 10^{10} L_{\odot}$). At higher redshifts ($0.5 \lesssim z \lesssim 1$), luminous infrared galaxies (LIRGs, $10^{11} L_{\odot} \leq L_{\text{IR}} \leq 10^{12} L_{\odot}$) become the dominant population among the sources detected at $24 \mu\text{m}$, while we also detect a significant number of more modest starbursts and spirals. The most extreme sources such as the ultra-luminous IR galaxies (ULIRGs, $L_{\text{IR}} \geq 10^{12} L_{\odot}$) still remain quite rare up to $z \sim 1.2$.

2. Evolution of the infrared luminosity function

Based on this sample of $24 \mu\text{m}$ -selected sources, and using the libraries of IR SEDs for computing the k -corrections, we derived monochromatic luminosity functions (LFs) at different IR wavelengths and in various redshift bins up to $z \sim 1$. Figure 1 shows these luminosity functions plotted at $60 \mu\text{m}$ and compared to the local $60 \mu\text{m}$ IRAS LF from Takeuchi et al. (2003). A strong evolution is clearly noticeable. We find that it can not be described by a uniform increase of the density of the local IR galaxy population (“pure density” evolution), but requires a shift of the characteristic luminosity L^* by a factor of $(1+z)^{4.0 \pm 0.5}$. Note that the well-known degeneracy between a pure evolution in luminosity or an evolution combining an increase in both density and luminosity can not be broken at this stage.




Figure 1. Luminosity functions estimated at $60 \mu\text{m}$ with the $1/V_{\text{max}}$ formalism in different redshift bins between $z=0$ and $z=1.2$ (* symbols). The 3σ uncertainties are indicated with vertical solid lines. Data points can be fitted by a pure luminosity evolution ($L_* \propto (1+z)^{4.2}$, solid curve) or a combination of luminosity and density evolution ($L_* \propto (1+z)^{3.5}$, $\Phi_* \propto (1+z)^{1.0}$, dashed curve) of the local $60 \mu\text{m}$ luminosity function (dotted curve). Vertical dashed-dotted lines correspond to the 80% completeness limit in each redshift bin.

3. Star formation history up to $z \sim 1$

Star-forming galaxies are believed to be the major component of the $24 \mu\text{m}$ source population (Silva et al. 2004). Assuming the calibration between the star formation rate of galaxies and their infrared luminosities (Kennicutt 1998), we converted the evolution of the luminosity functions into an history of the star formation up to $z \sim 1$. Results are shown on Figure 2 where the relative contributions of normal starbursts, LIRGs and ULIRGs are also illustrated. We find that IR luminous systems ($L_{\text{IR}} \geq 10^{11} L_{\odot}$) become the dominant population contributing to the comoving IR energy density beyond $z \sim 0.5-0.6$ and represent $\sim 70\%$ of the star-forming activity at $z \sim 1$. These preliminary results will be further developed by Le Floc'h et al. (2005, in prep.).

References

Blain, A., Smail, I., Ivison, R., Kneib, J.-P., & Frayer, D. 2002, Physics Reports, 369, 111




Figure 2. The star formation history up to $z \sim 1.2$ as seen by MIPS $24 \mu\text{m}$ (thick solid line), decomposed into the contributions of normal galaxies ($L_{\text{IR}} \leq 10^{11} L_{\odot}$), LIRGs and ULIRGs (shaded regions). Uncertainties on these estimates are still significant and they are not reported for clarity. Filled squares correspond to the contribution of *detected* MIPS sources, while empty diamonds and the empty square are from Flores et al. 1999 and Thompson et al. 2001, respectively. Model predictions from Chary & Elbaz (2001, cross-hatched region), Blain et al. (2002, dotted line) and Lagache et al. (2004, dashed line) are also plotted.

- Chapman, S. C., Helou, G., Lewis, G. F., & Dale, D. A. 2003, *ApJ*, 588, 186
 Chary, R. & Elbaz, D. 2001, *ApJ*, 556, 562
 Dale, D. A., Helou, G., Contursi, A., Silbermann, N. A., & Kolhatkar, S. 2001, *ApJ*, 549, 215
 Elbaz, D., Cesarsky, C. J., Fadda, D., et al. 1999, *A&A*, 351, L37
 Flores, H., Hammer, F., Désert, F. X., et al. 1999, *A&A*, 343, 389
 Kennicutt, R. C. 1998, *ARA&A*, 36, 189
 Lagache, G., Dole, H., & Puget, J.-L. 2003, *MNRAS*, 338, 555
 Le Fèvre, O. L., Vettolani, G., Paltani, S., et al. 2004, submitted to *A&A* (astro-ph/0403628)
 Silva, L., Maiolino, R., & G.L., G. 2004, *MNRAS*, in press (astro-ph/0403381)
 Smail, I., Ivison, R. J., & Blain, A. W. 1997, *ApJ*, 490, L5
 Szokoly, G. P., Bergeron, J., Hasinger, G., et al. 2004, *ApJS*, in press (astro-ph/0312324)
 Takeuchi, T. T., Yoshikawa, K., & Ishii, T. T. 2003, *ApJ*, 587, L89
 Thompson, R. I., Weymann, R. J., & Storrie-Lombardi, L. J. 2001, *ApJ*, 546, 694
 Vanzella, E., Cristiani, S., Dickinson, M., et al. 2004, submitted to *A&A* (astro-ph/0406591)
 Wolf, C., Meisenheimer, K., Kleinheinrich, M., et al. 2004, *A&A*, 421, 913